

SEARCH REQUEST FORM

4.381

Requestor's
Name:

Bottino

Serial

Number:

06/315,882

Date:

4/22/96

Phone:

305-6335

Art Unit:


1202

Search Topic:

Please write a detailed statement of search topic. Describe specifically as possible the subject matter to be searched. Define any terms that may have a special meaning. Give examples or relevant citations, authors keywords, etc., if known. For sequences, please attach a copy of the sequence. You may include a copy of the broadest and/or most relevant claim(s).

John Please

Possible
R's in A, B, C
ing (see p 3)



STAFF USE ONLY

Date completed:

4-25-96

Searcher:

JOHN DANTZMAN

Terminal time:

12 16

Elapsed time:

CPU time:

Total time:

26

Number of Searches:

5

Number of Databases:

5

Search Site

STIC

✓ CM-1

Pre-S

Type of Search

N.A. Sequence

A.A. Sequence

✓ Structure

Bibliographic

Vendors

IG Suite

✓ STN

Dialog

APS

Geninfo

SDC

DARC/Questel

Other

Reg Butst Manpat

=> FIL REG

FILE 'REGISTRY' ENTERED AT 17:28:07 ON 25 APR 96
 USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
 COPYRIGHT (C) 1996 American Chemical Society (ACS)

STRUCTURE FILE UPDATES: 19 APR 96 HIGHEST RN 175414-60-5
 DICTIONARY FILE UPDATES: 24 APR 96 HIGHEST RN 175414-60-5


TSCA INFORMATION NOW CURRENT THROUGH DECEMBER 1995

Please note that search-term pricing does apply when
 conducting SmartSELECT searches.

=>

=> D QUE L10

L5 STR



CH2=C
 26 @27

O~C
 29 @30

NC~C
 32 @33

O2N~C
 35 @36

VAR G1=27/30/33/36

VAR G2=19/20/21/22

NODE ATTRIBUTES:

CONNECT IS E1 RC AT 29

DEFAULT MLEVEL IS ATOM

DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED


NUMBER OF NODES IS 32

STEREO ATTRIBUTES: NONE

L7 129 SEA FILE=REGISTRY SSS FUL L5

L9 STR

parent



CH2=C	O~C	NC~C	O2N~C
26 @27	29 @30	32 @33	35 @36

VAR G1=27/30/33/36

VAR G2=19/20/21/22

NODE ATTRIBUTES:

CONNECT IS E1 RC AT 29

DEFAULT MLEVEL IS ATOM

DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED

NUMBER OF NODES IS 32

STEREO ATTRIBUTES: NONE

L10 16 SEA FILE=REGISTRY SUB=L7 SSS FUL L9

=>

=> FIL CAPLUS

FILE 'CAPLUS' ENTERED AT 17:28:17 ON 25 APR 96
USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
COPYRIGHT (C) 1996 AMERICAN CHEMICAL SOCIETY (ACS)

FILE COVERS 1967 - 25 Apr 1996 VOL 124 ISS 18
FILE LAST UPDATED: 25 Apr 1996 (960425/ED)

To help control your online searching costs, consider using the
HCAPLUS file when using the FSEARCH command or when conducting
SmartSELECT searches with large numbers of terms.

A new table-of-contents alerting feature is available in the
CAPLUS file. See NEWS or enter HELP TOC for details.

Thesauri are now available for the WIPO International Patent
Classifications (IPC) editions 1-6 in the /IC1, /IC2, /IC3, /IC4,
/IC5, and /IC (/IC6) fields, respectively. The thesauri in the
/IC5 and /IC fields also include the corresponding catchword terms
from the IPC subject headings and subheadings.


=> S L10

L11 5 L10

=> D BIB ABS HITSTR

L11 ANSWER 1 OF 5 CAPLUS COPYRIGHT 1996 ACS
AN 1995:874329 CAPLUS
DN 123:329474
TI Active-site conformation of 17-(3-pyridyl)androsta-5,16-dien-3.beta.-ol, a potent inhibitor of the P450 enzyme C17.alpha.-hydroxylase/C17-20 lyase
AU Burke, David F.; Laughton, Charles A.; Snook, Chris F.; Neidle, Stephen
CS Cancer Research Campaign Biomolecular Structure Unit, Institute Cancer Research, Sutton, Surrey, SM2 5NG, UK
SO Bioorg. Med. Chem. Lett. (1995), 5(11), 1125-30
CODEN: BMCLE8; ISSN: 0960-894X
DT Journal
LA English
AB 17-(3-Pyridyl)androsta-5,16-dien-3.beta.-ol, a nanomolar inhibitor of the P 450 enzyme C17.alpha.-hydroxylase/C17-20 lyase, is a target for prostate cancer chemotherapy. A model is presented for the inhibitor docked into the structure of the enzyme.
IT 154229-19-3
RL: BPR (Biological process); BIOL (Biological study); PROC (Process)
(active-site conformation of 17-(3-pyridyl)androsta-5,16-dien-3.beta.-ol, a potent inhibitor of the P 450 enzyme C17.alpha.-hydroxylase/C17-20 lyase)
RN 154229-19-3 CAPLUS
CN Androsta-5,16-dien-3-ol, 17-(3-pyridinyl)-, (3.beta.)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



=> D BIB ABS HITSTR 2

L11 ANSWER 2 OF 5 CAPLUS COPYRIGHT 1996 ACS
AN 1995:723267 CAPLUS
DN 123:112515
TI Synthesis of 17-(3-pyridyl) steroids
IN Potter, Gerard Andrew; Hardcastle, Ian Robert
PA British Technology Group Ltd., UK
SO Brit. UK Pat. Appl., 17 pp.
CODEN: BAXXDU
PI GB 2282377 A1 950405
AI GB 94-19139 940922
PRAI GB 93-20132 930930
GB 94-14192 940714
DT Patent
LA English
OS CASREACT 123:112515; MARPAT 123:112515
AB 17-(3-Pyridinyl)-substituted steroids are prepd. by subjecting a 17-iodo or -bromo steroid to a palladium complex-catalyzed cross-coupling reaction with a (3-pyridyl)-substituted borane in a proportion of at least 1.0 equiv. of borane per equiv. of steroid, in an org. solvent, and optionally esterifying the resulting 3.beta.-hydroxy steroid. Thus, dehydroepiandrosterone was converted to its hydrazone and then to its iodide. The latter compd. was treated with 1.1 equiv. diethyl(3-pyridyl)borane and then acetylated to give 3.beta.-acetoxy-17-(3-pyridyl)androsta-5,16-diene.
IT 154229-19-3P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation) (synthesis of 17-(3-pyridyl) steroids)
RN 154229-19-3 CAPLUS
CN Androsta-5,16-dien-3-ol, 17-(3-pyridinyl)-, (3.beta.)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



=> D BIB ABS HITSTR 3

L11 ANSWER 3 OF 5 CAPLUS COPYRIGHT 1996 ACS
AN 1995:608136 CAPLUS
DN 123:83811
TI Novel Steroidal Inhibitors of Human Cytochrome P45017.alpha.-
Hydroxylase-C17,20-lyase): Potential Agents for the Treatment of
Prostatic Cancer
AU Potter, Gerard A.; Barrie, S. Elaine; Jarman, Michael; Rowlands,
Martin G.
CS Cancer Research Campaign Centre for Cancer Therapeutics, Institute
of Cancer Research, Sutton/Surrey, SM2 5NG, UK
SO J. Med. Chem. (1995), 38(13), 2463-71
CODEN: JMCMAR; ISSN: 0022-2623
DT Journal
LA English
OS CJACS-IMAGE; CJACS
AB Steroidal compds. having a 17-(3-pyridyl) substituent together with
a 16,17-double bond have been synthesized using a
palladium-catalyzed cross-coupling reaction of a 17-enol triflate
with diethyl(3-pyridyl)borane and are potent inhibitors of human
testicular 17.alpha.-hydroxylase-C17,20-lyase. The requirement for
these structural features is stringent: compds. having 2-pyridyl,
4-pyridyl, or 2-pyridylmethyl substituents instead of 3-pyridyl
substituents were either poor inhibitors or noninhibitory. Redn. of
the 16,17-double bond to give 17.beta.-pyridyl derivs. diminished
potency with 3-pyridyl substitution, but increased it with a
4-pyridyl substituent present. In contrast, a variety of
substitution patterns in rings A-C of the steroid skeleton was
tolerated. The most potent compds. are candidates for development
as drugs for the treatment of hormone-dependent prostatic carcinoma.
IT 154229-19-3P 154229-25-1P 165334-72-5P
RL: BAC (Biological activity or effector, except adverse); RCT
(Reactant); SPN (Synthetic preparation); BIOL (Biological study);
PREP (Preparation)
(prepn. of novel steroidal inhibitors of human cytochrome P
45017.alpha.-hydroxylase-C17,20-lyase)
RN 154229-19-3 CAPLUS
CN Androsta-5,16-dien-3-ol, 17-(3-pyridinyl)-, (3.beta.)- (9CI) (CA
INDEX NAME)


Absolute stereochemistry.



RN 154229-25-1 CAPLUS

CN Androst-16-en-3-ol, 17-(3-pyridinyl)-, (3.alpha.,5.alpha.)- (9CI)
(CA INDEX NAME)


Absolute stereochemistry.



RN 165334-72-5 CAPLUS

CN Androsta-5,16-dien-3-ol, 17-(4-pyridinyl)-, (3.beta.)- (9CI) (CA
INDEX NAME)

Absolute stereochemistry.




IT 154229-21-7P 154229-23-9P 154229-26-2P
154229-27-3P 165334-71-4P

RL: BAC (Biological activity or effector, except adverse); SPN
(Synthetic preparation); BIOL (Biological study); PREP (Preparation)
(prepn. of novel steroidal inhibitors of human cytochrome P
45017.alpha.-hydroxylase-C17,20-lyase)

RN 154229-21-7 CAPLUS

CN Androsta-4,16-dien-3-one, 17-(3-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 154229-23-9 CAPLUS

CN Estra-1,3,5(10),16-tetraen-3-ol, 17-(3-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 154229-26-2 CAPLUS

CN Androst-16-en-3-one, 17-(3-pyridinyl)-, (5.alpha.)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 154229-27-3 CAPLUS

CN Androsta-4,16-diene-3,11-dione, 17-(3-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 165334-71-4 CAPLUS

CN Androsta-5,16-dien-3-ol, 17-(2-pyridinyl)-, (3.beta.)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



=> D BIB ABS HITSTR 4

L11 ANSWER 4 OF 5 CAPLUS COPYRIGHT 1996 ACS
AN 1994:645446 CAPLUS
DN 121:245446
TI Pharmacology of novel steroidal inhibitors of cytochrome
P45017.alpha. (17.alpha.-hydroxylase/C17-20 lyase)
AU Barrie, S. E.; Potter, G. A.; Goddard, P. M.; Haynes, B. P.;
Dowsett, M.; Jarman, M.
CS Drug Development Sec., Inst. Cancer Res., Sutton, SM2 5NG, UK
SO J. Steroid Biochem. Mol. Biol. (1994), 50(5-6), 267-73
CODEN: JSBBEZ; ISSN: 0960-0760
DT Journal
LA English
AB Medical or surgical castration for the treatment of prostatic
cancers prevents androgen prodn. by the testes, but not by the
adrenals. Inhibition. of the key enzyme for androgen biosynthesis,
cytochrome P 45017.alpha., could prevent androgen prodn. from both
sources. The in vivo effects of 17-(3-pyridyl)androsta-5,16-dien-
3.beta.-ol (CB7598) and 17-(3-pyridyl)androsta-5,16-dien-3-one
(CB7627), novel potent steroidal inhibitors of this enzyme, on WHT
mice were compared with those of castration and two clin. active
compds., ketoconazole and flutamide. Flutamide and surgical
castration caused significant redns. in the wts. of the ventral
prostate and seminal vesicles. CB7598, in its 3.beta.-O-acetate
form (CB7630), and CB7627 caused significant redns. in the wts. of
the ventral prostate, seminal vesicles, kidneys and testes when
administered once daily for 2 wks. Ketoconazole, given on the same
schedule, caused no redns. Plasma testosterone was reduced to
.ltoreq. 0.1 nM by CB7630, despite a 3- to 4-fold increase in the
plasma level of LH. Adrenal wts. were unchanged following treatment
with CB7630 or CB7627 but were markedly increased following
ketoconazole, indicating no inhibition. of corticosterone prodn. by
these steroidal compds. These results indicate that CB7598, CB7630
or CB7627 may be useful in the treatment of hormone-dependent
prostatic cancers.
IT 154229-19-3, CB 7598 154229-21-7, CB 7627
RL: BAC (Biological activity or effector, except adverse); THU
(Therapeutic use); BIOL (Biological study); USES (Uses)
(pharmacol. of steroidal inhibitors of cytochrome P 45017.alpha.
(17.alpha.-hydroxylase/C17-20 lyase))
RN 154229-19-3 CAPLUS
CN Androsta-5,16-dien-3-ol, 17-(3-pyridinyl)-, (3.beta.)- (9CI) (CA
INDEX NAME)


Absolute stereochemistry.



RN 154229-21-7 CAPLUS


CN Androsta-4,16-dien-3-one, 17-(3-pyridinyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



=> D BIB ABS HITSTR 5

L11 ANSWER 5 OF 5 CAPLUS COPYRIGHT 1996 ACS
 AN 1994:270958 CAPLUS
 DN 120:270958
 TI 17-(3-Pyridyl)-substituted steroids useful in cancer treatment
 IN Barrie, Susan Elaine; Jarman, Michael; Potter, Gerard Andrew
 PA British Technology Group Ltd., UK
 SO Brit. UK Pat. Appl., 40 pp.
 CODEN: BAXXDU
 PI GB 2265624 A1 931006
 AI GB 93-5269 930315
 PRAI GB 92-7057 920331
 GB 92-24880 921127
 DT Patent
 LA English
 OS MARPAT 120:270958
 GI



AB The title compds. are useful for treatment of androgen-dependent disorders, esp. prostatic cancer, and also estrogen-dependent disorders such as breast cancer. Claimed are compds. of formula I [X = steroid A-B-C ring residue; R = H, alkyl; R14 = H, halo, alkyl; R15 = H, alkyl, alkoxy, OH, alkylcarbonyloxy; or R15R15 = oxo, CH2; or R14R15 = pi bond and the other R15 = H, alkyl; R16 = H, halo, alkyl], in the form of free bases or pharmaceutically acceptable acid addn. salts, with the proviso that 5 specific compds. are claimed only for use in therapy. For example, dehydroepiandrosterone 3-acetate [i.e. 3.beta.-acetoxyandrost-5-en-17-one] was treated with (CF3SO2)2O and 2,6-di-tert-butyl-4-methylpyridine in CH2Cl2 to give 58% androsta-3,5,16-trien-17-yl trifluoromethanesulfonate. This triflate was coupled with diethyl(3-pyridyl)borane in the presence of Pd(PPh3)2Cl2 (84%), followed by hydrolytic deacetylation with aq. NaOH (79%), to give pyridylandrostadienol II. The IC50 values of II for inhibition of C17-C20 lyase and 17.alpha.-hydroxylase in vitro were, resp., 0.0029 .mu.M and 0.0040 .mu.M (cf. 0.026 and 0.065 for ketoconazole). In vivo organ wt. and endocrine test results for appropriate I in mice indicated inhibition of androgen synthesis, particularly testosterone.

IT 154229-19-3P 154229-21-7P 154229-23-9P

154229-25-1P 154229-26-2P 154229-27-3P

154229-29-5P 154229-30-8P 154229-33-1P


154229-34-2P 154229-35-3P 154229-47-7P

RL: SPN (Synthetic preparation); PREP (Preparation)
 (prepn. of, for treatment of hormone-dependent cancer)

RN 154229-19-3 CAPLUS

CN Androsta-5,16-dien-3-ol, 17-(3-pyridinyl)-, (3.beta.)- (9CI) (CA
 INDEX NAME)


Absolute stereochemistry.



RN 154229-21-7 CAPLUS

CN Androsta-4,16-dien-3-one, 17-(3-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 154229-23-9 CAPLUS

CN Estra-1,3,5(10),16-tetraen-3-ol, 17-(3-pyridinyl)- (9CI) (CA INDEX
 NAME)


Absolute stereochemistry.



RN 154229-25-1 CAPLUS

CN Androst-16-en-3-ol, 17-(3-pyridinyl)-, (3.alpha.,5.alpha.)- (9CI)
(CA INDEX NAME)


Absolute stereochemistry.



RN 154229-26-2 CAPLUS

CN Androst-16-en-3-one, 17-(3-pyridinyl)-, (5.alpha.)- (9CI) (CA INDEX
NAME)


Absolute stereochemistry.



RN 154229-27-3 CAPLUS

CN Androsta-4,16-diene-3,11-dione, 17-(3-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 154229-29-5 CAPLUS

CN Androsta-4,16-dien-3-one, 6-fluoro-17-(3-pyridinyl)-, (6.beta.)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 154229-30-8 CAPLUS

CN Androsta-4,16-dien-3-one, 6-fluoro-17-(3-pyridinyl)-, (6.alpha.)-
(9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 154229-33-1 CAPLUS

CN Androsta-4,16-diene-3,6-dione, 17-(3-pyridinyl)- (9CI) (CA INDEX
NAME)


Absolute stereochemistry.



RN 154229-34-2 CAPLUS

CN Androst-16-en-3-ol, 17-(3-pyridinyl)-3-(trifluoromethyl)-,
(3.beta.)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 154229-35-3 CAPLUS

CN Androsta-5,16-dien-3-ol, 17-(5-methyl-3-pyridinyl)-, (3.beta.)-
(9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 154229-47-7 CAPLUS

CN Androsta-3,5,16-trien-3-ol, 17-(3-pyridinyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 154229-45-5 154229-46-6


RL: RCT (Reactant)

(therapeutic use of)

RN 154229-45-5 CAPLUS

CN Androsta-5,14,16-trien-3-ol, 17-(3-pyridinyl)-, (3.beta.)- (9CI)
(CA INDEX NAME)


Absolute stereochemistry.



RN 154229-46-6 CAPLUS

CN Androsta-5,16-diene-3,15-diol, 17-(3-pyridinyl)-, 15-acetate,
(3.β.,15.β.)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



=> FIL CAOLD

FILE 'CAOLD' ENTERED AT 17:29:11 ON 25 APR 96
USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
COPYRIGHT (C) 1996 AMERICAN CHEMICAL SOCIETY (ACS)

FILE COVERS 1957-1966

FILE LAST UPDATED: 30 OCT 91 (910803/ED)

To help control your online searching costs, consider using the
HCAOLD File when conducting SmartSELECT searches with large
numbers of terms.

=> D HIS

(FILE 'HOME' ENTERED AT 17:21:55 ON 25 APR 96)

FILE 'REGISTRY' ENTERED AT 17:21:59 ON 25 APR 96

L1 STR
L2 25 S L1
L3 STR L1
L4 35 S L3
L5 STR L3
L6 6 S L5
L7 129 S L5 FUL

FILE 'CAPLUS' ENTERED AT 17:27:07 ON 25 APR 96

L8 19 S L7

FILE 'REGISTRY' ENTERED AT 17:27:14 ON 25 APR 96

L9 STR L5
L10 16 S L9 SSS FUL SUB=L7

FILE 'REGISTRY' ENTERED AT 17:28:07 ON 25 APR 96

FILE 'CAPLUS' ENTERED AT 17:28:17 ON 25 APR 96
L11 5 S L10

FILE 'CAOLD' ENTERED AT 17:29:11 ON 25 APR 96

L12 0 S L10

FILE 'BEILSTEIN' ENTERED AT 17:29:21 ON 25 APR 96

L13 0 S L9 FUL

FILE 'MARPAT' ENTERED AT 17:29:43 ON 25 APR 96



L14 6 S L7 FUL
L15 3 S L9 SSS FUL SUB=L14

=>

=>

=> D QUE L15

L5 STR



CH₂=C
26 @27

O~C
29 @30

NC~C
32 @33

O₂N~C
35 @36

VAR G1=27/30/33/36

VAR G2=19/20/21/22

NODE ATTRIBUTES:

CONNECT IS E1 RC AT 29

DEFAULT MLEVEL IS ATOM

DEFAULT ECLEVEL IS LIMITED



GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED

NUMBER OF NODES IS 32

STEREO ATTRIBUTES: NONE

L9 STR



CH₂=C
26 @27

O~C
29 @30

NC~C
32 @33

O₂N~C
35 @36

VAR G1=27/30/33/36

VAR G2=19/20/21/22

NODE ATTRIBUTES:

CONNECT IS E1 RC AT 29

DEFAULT MLEVEL IS ATOM

DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:
RING(S) ARE ISOLATED OR EMBEDDED
NUMBER OF NODES IS 32


STEREO ATTRIBUTES: NONE

L14 6 SEA FILE=MARPAT SSS FUL L5
L15 3 SEA FILE=MARPAT SUB=L14 SSS FUL L9

=> D QHIT BIB ABS

L15 ANSWER 1 OF 3 MARPAT COPYRIGHT 1996 ACS

MSTR 3




G1 = OH
MPL: disclosure

AN 123:112515 MARPAT
TI Synthesis of 17-(3-pyridyl) steroids
IN Potter, Gerard Andrew; Hardcastle, Ian Robert
PA British Technology Group Ltd., UK
SO Brit. UK Pat. Appl., 17 pp.
CODEN: BAXXDU
PI GB 2282377 A1 950405
AI GB 94-19139 940922
PRAI GB 93-20132 930930
GB 94-14192 940714
DT Patent
LA English
OS CASREACT 123:112515
AB 17-(3-Pyridinyl)-substituted steroids are prepd. by subjecting a 17-iodo or -bromo steroid to a palladium complex-catalyzed cross-coupling reaction with a (3-pyridyl)-substituted borane in a proportion of at least 1.0 equiv. of borane per equiv. of steroid, in an org. solvent, and optionally esterifying the resulting 3.beta.-hydroxy steroid. Thus, dehydroepiandrosterone was converted to its hydrazone and then to its iodide. The latter compd. was treated with 1.1 equiv. diethyl(3-pyridyl)borane and then acetylated to give 3.beta.-acetoxy-17-(3-pyridyl)androsta-5,16-diene.


=> D QHIT BIB ABS 2

L15 ANSWER 2 OF 3 MARPAT COPYRIGHT 1996 ACS

MSTR 1



G1 = 30-1 26-5



G7 = OH

DER: or steroid derivatives

MPL: claim 1

NTE: substitution is restricted

AN 120:270958 MARPAT

TI 17-(3-Pyridyl)-substituted steroids useful in cancer treatment

IN Barrie, Susan Elaine; Jarman, Michael; Potter, Gerard Andrew

PA British Technology Group Ltd., UK

SO Brit. UK Pat. Appl., 40 pp.

CODEN: BAXXDU

PI GB 2265624 A1 931006

AI GB 93-5269 930315


PRAI GB 92-7057 920331

GB 92-24880 921127




DT Patent

LA English



GI




AB The title compds. are useful for treatment of androgen-dependent disorders, esp. prostatic cancer, and also estrogen-dependent disorders such as breast cancer. Claimed are compds. of formula I [X = steroid A-B-C ring residue; R = H, alkyl; R14 = H, halo, alkyl; R15 = H, alkyl, alkoxy, OH, alkylcarbonyloxy; or R15R15 = oxo, CH2; or R14R15 = pi bond and the other R15 = H, alkyl; R16 = H, halo, alkyl], in the form of free bases or pharmaceutically acceptable acid addn. salts, with the proviso that 5 specific compds. are claimed only for use in therapy. For example, dehydroepiandrosterone 3-acetate [i.e. 3.beta.-acetoxyandrost-5-en-17-one] was treated with (CF₃SO₂)₂O and 2,6-di-tert-butyl-4-methylpyridine in CH₂Cl₂ to give 58% androsta-3,5,16-trien-17-yl trifluoromethanesulfonate. This triflate was coupled with diethyl(3-pyridyl)borane in the presence of Pd(PPh₃)₂Cl₂ (84%), followed by hydrolytic deacetylation with aq. NaOH (79%), to give pyridylandrostadienol II. The IC₅₀ values of II for inhibition of C17-C20 lyase and 17.alpha.-hydroxylase in vitro were, resp., 0.0029 .mu.M and 0.0040 .mu.M (cf. 0.026 and 0.065 for ketoconazole). In vivo organ wt. and endocrine test results for appropriate I in mice indicated inhibition of androgen synthesis, particularly testosterone.




G10 = OH / F / Cl / Br / I / alkoxy<(1-8)> /
 alkenyl<(2-10)> / 33 / 47 / SH / 50 / 52 /
 54) Q (0-) N (0-) O (0-) S (0) OTHERQ, RC (1),
 RS (1) M5 (1) X7> (SO) / 76 / 79 /
 Hy<EC (1-




G11 = Ph / naphthyl
 G12 = Ph / naphthylV / alkoxy<(1-8)> / alkenyl<(2-10)> /
 33 / 47 / SH / 50 / 52 / 54) Q (0-) N (0-) O (0-) S (0)
 OTHERQ, RC (1), RS (1) M5 (1) X7> (SO) / 76 / 79 /
 Hy<EC (1-







G13 = OH / alkoxy<(1-3)> / CN / 34 / NO2 / F / Cl / Br /
 I / N10)> / 33 / 47 / SH / 50 / 52 /
 54) Q (0-) N (0-) O (0-) S (0) OTHERQ, RC (1),
 RS (1) M5 (1) X7> (SO) / 76 / 79 /
 Hy<EC (1-




G14 = H / alkyl<(1-8)> (SO) / Ph (SO) / naphthyl (SO)
 G15 = Hy<EC (1-3) Q (0-) N (0-) O (0-) S (0) OTHERQ,
 RC (1), RS (1) M5 (1) X7> / Hy<EC (1-3) Q (0-) N (0-) O (0-)
 S (0) OTHERQ (6-) C, AR (1-), BD (6-) N, RC (2),
 RS (0-) E5 (1-) E6 (0-) E7 (0) OTHER>
 G16 = Hy<EC (1-3) Q (0-) N (0-) O (1-) S (0) OTHERQ,
 AN (1-) S, RC (1), RS (1) M5 (1) X7> /
 Hy<EC (1-3) Q (0-) N (0-) O (1-) S (0) OTHERQ (6-) C,
 AN (1-) S, AR (1-), BD (6-) N, RC (2),
 RS (0-) E5 (1-) E6 (0-) E7 (0) OTHER>
 G17 = Hy<EC (1-3) Q (0-) N (0-) O (0-) S (0) OTHERQ,
 RC (1), RS (1) M5 (1) X7> (SO) /
 Hy<EC (1-3) Q (0-) N (0-) O (0-) S (0) OTHERQ (6-) C,
 AR (1-), BD (6-) N, RC (2), RS (0-) E5 (1-) E6 (0-) E7 (0)
 OTHER> (SO)
 G18 = Hy<EC (1-3) Q (0-) N (0-) O (1-) S (0) OTHERQ,
 AN (1-) S, RC (1), RS (1) M5 (1) X7> (SO) /
 Hy<EC (1-3) Q (0-) N (0-) O (1-) S (0) OTHERQ (6-) C,
 AN (1-) S, AR (1-), BD (6-) N, RC (2),
 RS (0-) E5 (1-) E6 (0-) E7 (0) OTHER> (SO)
 G19 = Hy<EC (1-3) Q (0-) N (0-) O (1-) S (0) OTHERQ,
 AN (1-) S, RC (1), RS (1) M5 (1) X7> (SO) /
 Hy<EC (1-3) Q (0-) N (0-) O (1-) S (0) OTHERQ (6-) C,
 AN (1-) S, AR (1-), BD (6-) N, RC (2),
 RS (0-) E5 (1-) E6 (0-) E7 (0) OTHER>
 (SO)



G20 = S / S(O) / SO2
 G21 = H / alkyl<(1-8)> / CH2Ph / cyclohexyl
 G22 = O / S
 G23 = 132 / 90 / OH / OEt / 100 / NHPH / NH2 / 165



G24 = Ph / NO2 / NH2 / NHCOMe / CN / CONH2 / NMe2 / 149 /
 OMe / 175




G25 = 2-pyridyl / Ph
 G26 = H / Ph
 G27 = COMe / CH(OH)Me / Bu-t
 G28 = 91 / 1-adamantyl / Bu-i / CH₂CH₂OH

p-C₆H₄G27
 91

G29 = CN / NO₂ / CONH₂
 G30 = CN / CONH₂
 G1 +G2 = NULL
 G3 +G4 = NULL
 DER: or pharmaceutically acceptable salts or esters
 MPL: claim 1
 NTE: substitution is restricted

AN 120:245602 MARPAT
 TI Preparation of 17-ethers and thioethers of 4-aza-steroids as steroid reductase inhibitors
 IN Witzel, Bruce E.; Tolman, Richard L.; Rasmusson, Gary H.; Bakshi, Raman K.; Yang, Shu Shu
 PA Merck and Co., Inc., USA
 SO PCT Int. Appl., 68 pp.
 CODEN: PIXXD2
 PI WO 9323040 A1 931125
 DS W: AU, BB, BG, BR, CA, CZ, FI, HU, JP, KR, KZ, LK, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SK, UA, US
 RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, DE, DK, ES, FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG
 AI WO 93-US4746 930519
 PRAI US 92-886031 920520
 DT Patent
 LA English
 GI



I

AB Title compds. [I; a, b both = single bonds, and R2 = H; or a = double bond, b = single bond, and R2 = H; or a = single bond, b = double bond, and R2 = null; R1 = H, aryl, (aryl)alkyl; R3 = H, Me, Et, OH, NH2, SMe; R4 = (substituted) alkyl, aryl, heterocyclyl; Z = XR4, (CHR1)nXR4; X = O, S, SO, SO2], were prepd. as inhibitors of steroid 5.alpha.-reductase enzymes 1 and 2 (no data). The compds. are useful for the treatment of hyperandrogenic disease conditions and diseases of the skin and scalp. Thus, 17-hydroxymethyl-4-methyl-5.alpha.-4-azaandrostan-3-one and diphenyldiazomethane in CH2Cl2 were treated dropwise with BF3.Et2O to give 17-diphenylmethoxymethyl-4-methyl-5.alpha.-4-azaandrostan-3-one.

=> D HIS

(FILE 'HOME' ENTERED AT 17:21:55 ON 25 APR 96)

FILE 'REGISTRY' ENTERED AT 17:21:59 ON 25 APR 96

L1 STR
L2 25 S L1
L3 STR L1
L4 35 S L3
L5 STR L3
L6 6 S L5
L7 129 S L5 FUL

FILE 'CAPLUS' ENTERED AT 17:27:07 ON 25 APR 96

L8 19 S L7

FILE 'REGISTRY' ENTERED AT 17:27:14 ON 25 APR 96

L9 STR L5
L10 16 S L9 SSS FUL SUB=L7

FILE 'REGISTRY' ENTERED AT 17:28:07 ON 25 APR 96

FILE 'CAPLUS' ENTERED AT 17:28:17 ON 25 APR 96
L11 5 S L10

FILE 'CAOLD' ENTERED AT 17:29:11 ON 25 APR 96

L12 0 S L10

FILE 'BEILSTEIN' ENTERED AT 17:29:21 ON 25 APR 96

L13 0 S L9 FUL

FILE 'MARPAT' ENTERED AT 17:29:43 ON 25 APR 96

L14 6 S L7 FUL
L15 3 S L9 SSS FUL SUB=L14

=>